

## ENHANCEMENT OF KIDNEY AND LIVER RESPIRATORY ACTIVITY BY QUERCETIN SULFONATES IN RATS CHRONICALLY EXPOSED TO AMMONIUM FLUORIDE

Z Juzyszyn,<sup>a</sup> B Czerny, Z Myśliwiec, A Put  
Szczecin, Poland

**Summary:** The respiratory activity of liver and kidney slices of rats chronically exposed to ammonium fluoride was studied. It was found that a mixture of quercetin sulfonates stimulated tissue metabolism and exerted a protective effect in NH<sub>4</sub>F intoxication through normalization of respiratory activity.

**Keywords:** Ammonium fluoride, Fluoride in rats, Kidney respiration, Liver respiration, Quercetin sulfonates.

### INTRODUCTION

Fluorides in the atmosphere are the most hazardous pollutants when and where their concentrations exceed admittedly very low levels. Their high toxicity and continuous build-up in the environment have been a growing threat to mankind.<sup>1-3</sup>

One of the most serious effects of fluorine intoxication is impairment of the energy system of the organism, affecting both glycolysis and oxidative processes of cells.<sup>2,4</sup> Morphological, histochemical, and biochemical studies on the toxicity of fluorine compounds have revealed lesions to parenchymatous organs like kidneys and liver.<sup>3,5,6-9</sup>

Our previous work demonstrated a beneficial effect of quercetin on disturbances of lipid metabolism.<sup>5</sup> A protective effect of pollen (*Cernitin*) on the respiratory activity of rats tissues in ammonium fluoride intoxication was also proven.<sup>10</sup> We have now decided to continue the search for easily accessible and safe pharmaceuticals effective in reducing the toxic effects of fluorides.

### MATERIAL AND METHODS

The experiment was performed on 96 male Wistar rats with an initial body weight of ca. 300 g. The animals were exposed in a 1-m<sup>3</sup> toxicological chamber to ammonium fluoride, NH<sub>4</sub>F, as an aerosol at an average concentration of 2 mg F/m<sup>3</sup> (twice the work-period MAC) for 3 months (13 weeks) or 6 months (26 weeks) for 6 hr/day for 5 days each week. The rats were randomly divided into 12 groups with 6 to 10 rats per group that were treated as shown in the following Table.


---

<sup>a</sup>For Correspondence: Dr Zygmunt Juzyszyn, Department of Experimental and Clinical Pharmacology, Pomeranian Academy of Medicine, al. Powstańców Wlkp 72, 70-111 Szczecin, Poland. E-mail: zjuzyszyn@wp.pl

**Table.** Treatment protocol for the rat experiments

Groups		Treatment
3 Mo.	6 Mo.	
IA	IB	Non-exposed control
IIA	IIB	Quercetin 5 mg/kg bw/24 hr
IIIA	IIIB	Quercetin 20 mg/kg bw/24 hr
IVA	IVB	NH <sub>4</sub> F 2 mg/m <sup>3</sup> ( Exposed control)
VA	VB	NH <sub>4</sub> F 2 mg/m <sup>3</sup> + Quercetin 5 mg/kg bw/24 hr
VIA	VIB	NH <sub>4</sub> F 2 mg/m <sup>3</sup> + Quercetin 20 mg/kg bw/24 hr

A mixture containing 1:1 quercetin 5',8-disulfonic (Na<sub>2</sub>QDSA) and monosulfonic acid sodium salts (NaQSA-5'/NaQSA-8) synthesized at the Department of Inorganic Chemistry, Technical University of Rzeszów (Figure 1), was mixed with standard chow pellets containing 7 mg F/kg. Full consumption of quercetin pellets was ascertained before the animals were returned to normal chow. Rats had free access to tap water containing 0.2 mg F/L.



**Figure 1.** Quercetin 8,5-disulfonic acid sodium salts: NaQSA-8 (A) and NaQSA-5 (B).

As in our previous work,<sup>5</sup> an aerosol of NH<sub>4</sub>F was led into the 1-m<sup>3</sup> toxicological chamber by an atomizer inhaler (Ogarit, JV-S). Rats were kept at constant conditions: humidity (60%), temperature (20 deg C), air flow (10 m<sup>3</sup> /hr), and a 12/12-hr light/dark cycle. The concentration of fluoride ion in the chamber was measured once a week using an ion-selective fluorine electrode according to PN-83/Z-04093.07 norm. The average value from 12

measurements (3-month exposure period) deviated from 2 mg F/m<sup>3</sup> by 6.0%, with individual values ranging from 1.6 to 2.4 mg F/m<sup>3</sup>. The average value from 24 measurements (6-month exposure period) deviated from 2 mg F/m<sup>3</sup> by 4.8%, with individual values ranging from 1.6 to 2.5 mg F/m<sup>3</sup>. Operation of the fluorine dosing system was continuously monitored by checking the volume of NH<sub>4</sub>F solution fed to the chamber.


Rats were weighed every two weeks. At the end of the experiment the animals were lethally anaesthetized with ether. Liver and kidneys were removed, frozen immediately at -70°C, and stored for up to one month. Tissue respiration rates were monitored using Clark's oxygen electrode (5331 Standard Oxygen Probe, YSI Instrument Co., Ohio, USA) in 40-80 mg tissue slices suspended in 5 cm<sup>3</sup> of medium containing 50 mM KCl, 2 µM rotenone (inhibitor of complex I of the respiratory chain), and 50 mM phosphate buffer at pH 7.3 and 37°C. Respiration was started by adding 100 mg potassium succinate to the incubation chamber. This amount of respiratory substrate grossly exceeded the normal cellular level and served to maintain maximal respiratory rate.<sup>11</sup> Oxygen consumption was expressed in nanoatoms/hr/mg dry tissue. The initial concentration of oxygen was taken as 199 µM. Dry tissue mass was determined by drying for 5 hr at 100°C and weighing on a precision balance. Statistical analysis was done using Student's t-test for unpaired results and taking the level of significance as  $p < 0.05$ .

## RESULTS


A decrease of 10% in the body mass of rats exposed for 6 months to ammonium fluoride as compared with controls (group IVB *vs.* IB) was noted. Differences in body mass for controls receiving and not receiving quercetin (IA-B and IVA-B) were insignificant. Changes in oxygen consumption rates of liver and kidney tissue slices are given in Figures 2a and 2b, respectively.

Oxygen consumption rates of kidney slices taken from animals not exposed to NH<sub>4</sub>F were increased by quercetin sulfonates administered during 6 months. The increase was 76% at 5 and 39% at 20 mg/kg bw/24 hr (groups IIB and IIIB, respectively *vs.* IB) (Figure 2b).

After the rats were exposed to HN<sub>4</sub>F for 3 or 6 months, the following changes in tissue respiratory activity were observed: an exposure for 3 months led to a 29% increase in the respiratory activity in the liver slices (Figure 2a) and a 25% decrease in the kidney slices (Figure 2b). After 6 months, the respiratory activity of the liver tissue decreased by 22% (Figure 2a), while that of kidney tissue increased by 71% (Figure 2b). The differences were statistically significant, except for the decrease in the respiratory rate of liver after 6 months (groups IVA *vs.* IA and IVB *vs.* IB).



**Figure 2a.** Respiratory activity (control = 100%) of liver tissue of rats after 3 months (A groups) and after 6 months (B groups) of the experiment.



**Figure 2b.** Respiratory activity (control = 100%) of kidney tissue of rats after 3 months (A groups) and after 6 months (B groups) of the experiment.

\*Statistically significant ( $p < 0.05$ ) in comparison with the respective control Group IA or IB taken as 100%.

As also seen in Figures 2a and 2b, in the 3-month exposure groups, 5 mg/kg bw/24 hr quercetin sulfonates normalized the rate of oxygen consumption of both liver and kidney slices (groups VA vs. IA). Quercetin sulfonates at 20 mg/kg bw/24 hr normalized oxygen consumption of liver slices and decreased it by 24% in the case of kidney tissue (groups VIA vs. IA). When exposure to NH<sub>4</sub>F was for 6 months, quercetin sulfonates at 5 mg/kg bw/24 hr normalized the rate of oxygen consumption of liver slices and increased the kidney rate by 30% (groups VB vs. IB). The higher dose of quercetin sulfonates increased oxygen consumption of kidney slices by 56%. An insignificant increase of 22% was noted for the liver (groups VIB vs. IB).

### DISCUSSION

The aim of the present work was to study the effect of ammonium fluoride and a mixture of quercetin sulfonates on the respiratory activity of rat liver and kidney. Under conditions of the present experiment with maximal activation of succinate dehydrogenase, oxygen consumption rates can be regarded as an index of the metabolic activity of the cell.<sup>11</sup>

The overall condition of animals assessed on the basis of body weight deteriorated only in the case of exposure to NH<sub>4</sub>F for 6 months as evidenced by a 10% weight loss in relation to the control group (groups IVB vs. IB). This finding is in line with other reports on the toxic properties of fluorine compounds.<sup>1,2</sup> No significant changes in body weight were observed when animals exposed to NH<sub>4</sub>F were simultaneously given quercetin sulfonates. We are inclined to attribute this result to the protective action of quercetin. The weight of animals in the remaining groups did not change in comparison with the controls (groups IA and IB).

Changes in the respiratory activity of the liver and kidney clearly show that both organs play a complementary role in NH<sub>4</sub>F detoxification. The respiratory activity of liver tissue was increased after 3 months of exposure to NH<sub>4</sub>F and decreased after 6 months. A reverse pattern was observed for kidney tissue: respiratory rates were reduced after 3 months of exposure and increased after 6 months. Apparently, the role of kidneys becomes more important in prolonged exposure to ammonium fluoride.

When assessing the toxic effect of NH<sub>4</sub>F one has to remember that both components, *i.e.*, fluoride ion and ammonium ion may contribute to the properties of this compound. In present study NH<sub>4</sub><sup>+</sup> ion concentration applied as an aerosol was 1.9 mg/m<sup>3</sup>. There is an evidence, that NH<sub>4</sub><sup>+</sup> ion given as a NH<sub>4</sub>Cl salt inhibited tissue respiration in rats, but at higher concentrations than these used in the present study, *i.e.*, 0.6 g/kg bw.<sup>12</sup> No further data on effects of NH<sub>4</sub><sup>+</sup> ion on tissue respiration in animals chronically exposed to NH<sub>4</sub><sup>+</sup> are available.

Plant products containing quercetin have long been exploited to stimulate bile excretion, protect the liver, enhance renal filtration, and increase the excretion of xenobiotics. Furthermore, the toxicity of fluorine compounds has been traced to the stimulation of peroxidation of cellular lipids.<sup>13,14</sup> The antioxidative action of flavonoids like quercetin is due to the presence of hydroxyl groups, particularly in ring B (Figure 1), which may contain two (catechol) or three (pyrogallol) hydroxyls.<sup>15</sup> An additional feature of quercetin sulfonates used by us is their enhanced solubility in water without any loss of antioxidative properties and metal chelating strength.<sup>16</sup>

In the present work, either 5 or 20 mg/kg bw/24 hr of quercetin sulfonates given to rats during NH<sub>4</sub>F exposure (3 or 6 months) normalized the respiratory activity of liver tissue. Normalization of the respiratory activity of kidneys was observed only for the lower dose and shorter exposure period. The effect was particularly sharp in the case of kidney slices from rats given quercetin sulfonates (5 or 20 mg/kg bw/24 hr) during 6 months. We attribute these findings to stimulation by sulfonate derivatives of quercetin of cellular respiratory activity.

As already mentioned, the present work is a continuation of the search for safe and easily accessible pharmaceuticals that are effective in reducing the toxic effects of fluorides. In our previous study, a protective effect of quercetin sulfonates on hepatic metabolism of lipids in rats exposed to ammonium fluoride was observed. Quercetin salts normalized lipid content in serum, liver homogenates, and liver microsomal fraction in rats exposed to NH<sub>4</sub>F for 3 months.<sup>5</sup> The results of the present work confirm beneficial properties of quercetin salts. The mixture of quercetin sulfonates stimulated and normalized tissue respiratory activity and thus could be applied as a prophylactic remedy in fluoride intoxication of people who are exposed to excessive fluoride emission.

#### REFERENCES

- 1 Gumińska M. Biochemical mechanism of influence of fluorine on the living organism. *Folia Med Cracov* 1981;23:305-21.
- 2 Gumińska M. Wpływ fluorków na gospodarkę energetyczną *in vitro* i *in vivo* oraz związane z tym efekty biologiczne. [Influence of fluorides on *in vitro* and *in vivo* metabolism and related biological effects]. In: Machoy Z, editor. *Metabolism of Fluorine '94. Proceedings of the 6<sup>th</sup> Symposium on the Metabolism of Fluorine*; 1994 Sept. 15, Szczecin, Poland. Szczecin: Polskie Towarzystwo Biochemiczne & Szczecińskie Towarzystwo Naukowe; 1994. p 9-11.
- 3 Waldbott GL. Symposium on the non-skeletal phase of chronic fluorosis. *Fluoride* 1976;9:5-8.
- 4 Machoy Z. Wpływ związków fluoru na łańcuch oddechowy [The influence of fluorine compounds on the respiratory chain]. In: Machoy Z, editor. *Metabolism of Fluorine*. Warsaw, Poznań: PWN; 1982. p. 3-5.

- 5 Czerny B, Put A, Myśliwiec M, Juzyszyn Z. The influence of quercetin on some parameters of lipid metabolism in rats chronically exposed to ammonium fluoride. *Fluoride* 2000;33:27-32.
- 6 Kessabi M, Hamliri A, Braun JP, Rico AG. Experimental acute sodium fluoride poisoning in sheep: renal, hepatic and metabolic effects. *Fundam Appl Toxicol* 1985;5(6 Pt 1):1025-33.
- 7 Put A, Myśliwiec Z. Die Wirkung von Pollenextracten auf Experimenten mit Ammonium Fluorid hervorgerufene Leberveränderungen bei Ratten [The action of pollen extracts in experimental changes induced in rat liver by ammonium fluoride]. *Arzteitschrift für Naturheilverfahren* 1993;34:680-6.
- 8 Put A, Myśliwiec Z. Der Einfluß von Pollenextracten auf biochemische Indikatoren der Leberfunktion im Verlauf einer chronischer Intoxikation mit Ammoniumfluorid bei Ratten. Teil 11 [The influence of pollen extracts on biochemical indicators of liver function in rats chronically intoxicated with ammonium fluoride. Part 11]. *Arzteitschrift für Naturheilverfahren* 1994;35:351-5.
- 9 Singh M, Kanwar KS. Effect of fluoride on tissue enzyme activities in rat. *Biochemical and histochemical studies. Fluoride* 1981;14:132-41.
- 10 Juzyszyn Z, Put A, Myśliwiec Z, Ceglecka M, Ogoński, T, Gałka G. Influence of standardized pollen extracts (Cemitin) on the respiratory activity of some tissues of rats subchronically exposed to ammonium fluoride. *Phytomedicine* 1997;4(4):315-7.
- 11 Longmuir S, Bouke A. The measurement of the diffusion of oxygen through respiring tissue. *Biochem J* 1960;76:225-9.
- 12 Tupitska OM. Osoblivosti metabolichnykh protsesiv v tkanikakh vagitnikh tvaryn ta ikh plodiv pid vplivom NH<sub>4</sub>Cl [Metabolic processes in the tissues of pregnant animals and their fetuses affected by ammonium chloride]. *Ukr Biokhim Zh* 1997;69(2):41-5.
- 13 Shayiq RM, Raza H, Kidwai AM. Fluoride and lipid peroxidation: a comparative study in different rat tissues. *Bull Environ Contam Toxicol* 1986;37(1):70-6.
- 14 Soni MG, Kachole MS, Pawar SS. Alterations in drug metabolizing enzymes and lipid peroxidation in different rat tissues by fluoride. *Toxicol Lett* 1984;21(2):167-72.
- 15 Hu JP, Calomme M, Lasure A, de Bruyne T, Pieters L, Vlietinck A, Vanden Berghe DA. Structure-activity relationship of the flavonoids with superoxide scavenging activity. *Biol Trace Elem Res* 1995;47:327-31.
- 16 Kopacz M, Kopacz S. Sulfonowe pochodne niektórych flawonoidów i ich zastosowanie [Sulfonyl derivatives of some flavonoids and their applications]. *Proceedings of the 1<sup>st</sup> Conference on flavonoids and their applications; 1996; Rzeszów, Poland. p. 57-70.*